

作用機制	普通名稱	銹 蟎	葉 蟎類	薊 馬類	盾 介殼 蟲類	軟 介殼 蟲類	粉 介殼 蟲類	柑 桔 木 蝨	柑 桔 刺 粉 蝨	蚜 蟲類	椿 象 類	柑 桔 潛 葉 蛾	黑 點 刺 蛾	斜 紋 夜 蛾	毒 蛾 類	鳳 蝶	鱗 翅 目 害 蟲	東 方 果 實 蠅
10A + 3A	合賽芬普寧		◎															
10A + 12B	佈賜芬蟎		◎															
10B	依殺蟎		◎															
12A	汰芬隆	◎	○											○				
12B	芬佈賜	◎	◎															
12C	毆蟎多		◎															
13	克凡派			◎										○	○			
14	培丹									○		◎		○				
15	二福隆	◎	◎											○	○		○	
15 + 1A	福隆納乃得	◎										◎						
16	布芬淨				◎	○												
18	得芬諾													◎	○			
19	三亞蟎		◎															
20B	亞醜蟎		◎															
21A	芬殺蟎		◎															
	芬普蟎	◎	◎															
	得芬瑞		◎															
	畢汰芬		◎															
	畢達本	◎	◎															
21A + 12C	芬普毆蟎多		◎															
23	賜派芬		◎															
	賜派滅			○	◎	◎	◎											
25A	賽芬蟎		◎															
UN	新殺蟎	◎	◎															
UN	蟎離丹		◎															
	礦物油		◎		◎	◎	◎											
	斜紋夜蛾費洛蒙													◎				
methyl eugenol	含毒甲基丁香油																	◎
C5	扶吉胺		◎															
M2	可濕性硫黃	◎																
M2	石灰硫黃	◎																

◎ 代表已登記在該使用範圍。

○ 該藥劑已於國內登記於防治其他作物之相同或同群害蟲種類。

上述資料如有錯誤，以主管機關公告為準。